

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY**

PASTORAL / AG™

CYROLAB™ **SPRAY-ON**

BLOWFLY TREATMENT FOR SHEEP

ACTIVE CONSTITUENT: 60 g/L CYROMAZINE

For the protection for up to eleven weeks against strike by
cyromazine-susceptible strains of blowflies (*Lucilia cuprina*)
when applied by spraying on sheep with long wool.

Resistance may develop to any
chemical.



20 L

DIRECTIONS FOR USE

Restraints:

DO NOT USE on sheep which are producing or may in future produce milk that may be used or processed for human consumption.

Contraindications:

DO NOT USE on open wounds, e.g. those resulting from marking or mulesing operations.

Dosage and Administration:


Critical Use Comments:

For use on sheep more than six weeks after shearing.

Reduced effectiveness is likely if used on sheep with lumpy wool.

When used on sheep with less than 6 weeks wool a reduced period of protection may result.

The product should not be applied during gusty or windy weather as spray drift will reduce the period of protection.

MONTHS SINCE SHEARING	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NUMBER OF BANDS	NUMBER OF SHEEP TREATED PER 20 L	COMMENTS
FOR THE PREVENTION OF BODY STRIKE					<ol style="list-style-type: none">1. Use only a spray-on applicator recommended by the manufacturer.2. Ensure the applicator gun is clean.3. Calibrate applicator gun using an accurate volume measure.4. Ensure all areas to be protected are treated.5. When preventing body strike ensure bands overlap along the midline – see diagram.6. This product is not ideal for treating existing strikes. Use a diazinon knockdown product as a dressing for existing strikes.7. Heavy rain following the application of this product could diminish the period of protection.8. PASTORAL AG Cyrolab Spray-On is coloured for band identification. This scourable dye will disappear shortly after treatment. 
Up to 6	34	17	2	588	
7	38	19	2	526	
8	44	22	2	454	
9	50	25	2	400	
10	56	28	2	357	
Lambs less than 15 kg	17	17	1	1176	
FOR THE PREVENTION OF BODY STRIKE AND CRUTCH STRIKE					<ol style="list-style-type: none">9. PASTORAL AG Cyrolab Spray-On is coloured for band identification. This scourable dye will disappear shortly after treatment.
Up to 6	51	17	3	392	
7	57	19	3	350	
8	66	22	3	303	
9	75	25	3	266	
10	84	28	3	238	
Lambs less than 15 kg	34	17	2	588	
FOR THE PREVENTION OF POLL STRIKE					Rams and sheep susceptible to poll strike should be treated. Apply at least 5 mL of PASTORAL AG Cyrolab Spray-On to the areas requiring protection, around the poll and horns.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

General Directions:

DO NOT DILUTE. DO NOT mix with any other product.

Apply directly from the container using a spray-on applicator gun recommended by the manufacturer. Except for small lambs under 15 kg, the dose volume is determined by the number of months since shearing. Apply PASTORAL AG Cyrolab Spray-On according to the diagrams and tables in the Directions for Use. PASTORAL AG Cyrolab Spray-On may be applied to all breeds of sheep, providing protection against fly strike. PASTORAL AG Cyrolab Spray-On is applied as a low-volume treatment onto or near areas where protection is required. When applying, the nozzle of the applicator gun should be approximately 25 cm above the fleece, to achieve a 15 cm wide band at each pass.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 7 days before slaughter for human consumption.

MILK: DO NOT USE on sheep which are producing or may in future produce milk that may be used or processed for human consumption.

WOOL: DO NOT USE less than 2 months before shearing or fibre collection.

TRADE ADVICE – EXPORT SLAUGHTER INTERVAL (ESI)

DO NOT USE less than 28 days before slaughter for export. Before using this product, confirm the current ESI from the distributor on (03) 5820 8444 or the APVMA website (www.apvma.gov.au/residues).

Use of this product may result in wool residues that may not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ADDITIONAL USER SAFETY INFORMATION

Sheep Rehhandling Interval: After treatment wait until sheep are dry before rehhandling.

Additional information is listed in the safety data sheet.

ENVIRONMENTAL PROTECTION

DO NOT contaminate dams, rivers or streams with the chemical, discarded sump solution or used containers.

DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

KEEP OUT OF REACH OF CHILDREN. Store below 30°C (room temperature). Store in original container, tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container.

B:

EXP:

