PASTORAL AG

DORAMECTIN

Pour On Endectocide for Cattle





ACTIVE CONSTITUENT: 5mg/mL Doramectin

For the treatment and control of doramectin sensitive internal and external parasites of cattle. For the treatment and persistent control of gastrointestinal roundworms, lungworms and buffalo fly in cattle. For the control of eye worms, sucking and biting lice, mange mites and cattle ticks.

Persistent protection against re-infection of cattle by:
Small Brown Stomach Worm (Ostertagia ostertagi –35 days)
Small Intestinal Worm (Cooperia spp – 35 days)
Barbers Pole worm (Haemonchus placei 35 days)
Stomach Hair worm (Trichostrongylus axei 35 days)
Lung worm (Dictyocaulus viviparous 28 days)
When applied as directed, this product can persistently protect against various gastrointestinal roundworms for 21 to 35 days; against lungworms for up to 28 days and against buffalo flies for up to 14 days.
Pour-on Applicator Required.

DOSE RATE: 1ml/10kg bodyv

1ml/10kg bodyweight.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 42 days before slaughter for human consumption. **MILK:** Zero (0) days.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI):

DO NOT USE less than 42 days before slaughter for export.

APVMA Approval No.: 86118/121969

*See product label for full persistent protection claims.

APVMA Approval No.: 86118/121969



