

PASTORAL / AG[®]

IMIDALAB[™]

POUR-ON

LICE TREATMENT FOR SHEEP

ACTIVE CONSTITUENT: 35 g/L IMIDACLOPRID
Also contains: 300 g/L N-METHYL-2-PYRROLIDONE (solvent)

For the control of susceptible body lice *Bovicola ovis* on shorn sheep up to 7 days off shears and unshorn lambs up to 2 months of age. This product can be expected to protect sheep against re-infestation with lice for 4 weeks after treatment when applied within 24 hours of shearing.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 21 days before slaughter for human consumption.

MILK: DO NOT USE on ewes which are producing, or may in the future produce milk or milk products for human consumption.

WOOL: DO NOT USE less than 6 months before shearing or fibre collection.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI)
DO NOT slaughter for export less than 63 days after treatment.

RECOMMENDED APPLICATOR

Pastoral Ag 30mL Variable Applicator (or other suitable applicator) with 6 hole T Bar nozzle.

ADDITIONAL INFORMATION

To prevent re-infestation treated sheep must not be mixed with lice free sheep until 6 weeks after treatment. Live lice on ewes treated within 6 weeks of lambing can infect lambs from these ewes.

Sheep must be treated within 7 days of shearing. Unshorn lambs can be treated up to 2 months of age. Efficacy is not affected by moderate rainfall either before or after treatment (50mm rain 1 hour prior to treatment and/ or 12.5mm 2 hours after treatment and/or 25mm 24 hours after treatment). Treated sheep must not be mixed with lousy sheep.

DO NOT use on sheep which have not been cleanly shorn for any reason e.g. mycotic dermatitis.

Management practices which promote good lice control such as complete musters, accurate treatment of every animal according to label, ensuring all animals can be treated at the same time, no split shearings, stock-proof fencing to prevent external introductions,

quarantine and lice treatment of all newly purchased stock are essential for effective lice control.

Treated sheep could potentially develop backline dermatitis following prolonged exposure to extreme heat and sunlight without access to shade.



DO NOT leave container, hose or applicator in the sun for long periods of time.

Pastoral Ag Imidacloprid may be used in conjunction with worm control drenches and at the same time as vaccination.

Pastoral Ag Imidacloprid contains a blue dye to identify treated sheep. This dye is scourable from the wool. The product can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces.

Pastoral Ag Imidacloprid is safe for use on breeding rams.

DOSAGE TABLE

BODYWEIGHT (KG)	DOSE (ML)	METHOD	APPLY AS BELOW
Below 6kg	8	Single Stripe	
6 – 8	12		
8.1 – 10	15		
10.1 – 12.5	20		
12.6 – 15	25		
15.1 – 20	30		
20.1 – 30	40	Double Stripe	
30.1 – 55	60		
55.1 – 80	80		
80.1 – 100	90		
Add 10mL for each 10kg above 100kg			

AVAILABLE SIZES
20 L

For more information, visit pastoralag.com.au

For product enquiries, email us at enquiries@pastoralag.com.au or call us on (03) 5820 8444


APVMA Approval No.: 90295/128455



NEW
PACKAGING!

BOOST YOUR
LICE CONTROL

PROTECT AGAINST
REINFESTATION
FOR UP TO 4 WEEKS



EXCLUSIVELY AVAILABLE FROM

Elders

MADE IN AUSTRALIA