

PASTORAL / AG[®]

STRIKELAB[™]

SPRAY-ON SHEEP BLOWFLY TREATMENT

ACTIVE CONSTITUENT: 50 g/L DICYCLANIL

BLOCK THE STRIKE

**SEASON-LONG
PROTECTION!**

UP TO 24 WEEKS

Low volume treatment

**Suitable for all breeds
and any length of wool**

For the protection of sheep, either off-shears or with any length wool, against fly strike (*Lucilia cuprina*) for 18 to 24 weeks.

For the protection of mulesing and marking wounds on sheep against fly strike (*Lucilia cuprina*) during the wound healing process.

DOSING & ADMINISTRATION

Shake well before use.

Pastoral Ag Strikelab is applied as a low-volume treatment on to areas where protection is required. When applying Pastoral Ag Strikelab, the nozzle of the applicator gun should be high enough above the fleece (approximately 15 - 25 cm), to achieve a 15 cm wide band at each pass for protection of body strike. For treatment of mulesing wounds, the band must be wide enough to cover the wound and 25 mm onto the surrounding wool.

DO NOT apply during gusty or windy weather as spray drift may reduce the period of protection. It is good agricultural practice for mulesing/ marking to be carried out at times of minimal fly activity. Reduced efficacy may occur in sheep affected with dermatophilosis.

GENERAL DIRECTIONS

Pastoral Ag Strikelab provides season long protection to sheep against body, breach and poll strike and may be used either off-shears or on sheep with any length wool. Pastoral Ag Strikelab

also protects mulesing and marking wounds on lambs against fly strike during the wound healing process. The dose volume is determined by the weight of sheep being treated. Apply Pastoral Ag Strikelab according to the diagrams and tables in the Directions for Use. DO NOT DILUTE. Apply directly from the container or an appropriate back pack using a recommended spray-on applicator gun. Pastoral Ag Strikelab may be applied to all breeds of sheep to obtain protection against fly strike.

Significant soiling with dags, mud, dust or urine may reduce efficacy. Where significant dags are present, a light crutching is indicated prior to treatment. Protection period may be reduced if sheep scour heavily following application.

WITHHOLDING PERIODS

MEAT: DO NOT USE

less than 28 days before slaughter for human consumption.

MILK: DO NOT USE on sheep which are producing or may in the future produce milk or milk products for human consumption.

WOOL HARVEST INTERVAL: DO NOT USE less than 3 months before shearing or fibre collection.

ESI: DO NOT use less than 120 days before slaughter for export.

AVAILABLE SIZES

20 L

For more information, visit pastoralag.com.au

For product enquiries, email us at enquiries@pastoralag.com.au or call us on (03) 5820 8444



EXCLUSIVELY AVAILABLE FROM



MADE IN AUSTRALIA

