

JUR-O

FOR CATTLE

ACTIVE CONSTITUENT: 5 g/L IVERMECTIN

Pastoral Ag Iverlab Pour-On for Cattle gastrointestinal nematodes (including inhibited immature Ostertagia ostertagi), lungworms (Dictyocaulus viviparus), eyeworms (Thelazia spp.), sucking and biting lice, chorioptic and sarcoptic mange mites, buffalo flies (Haematobia irritans exigua) and cattle ticks

APPLICATOR

Young stock: Pastoral Ag 30ml Variable Applicator - Pour-On Nozzle. Stock over 300kg: 65ml Pour-On Applicator

WITHHOLDING PERIODS

MEAT: 28 days. MILK: 0 days. ESI: 28 days

STORAGE

Store below 30°C (room temperature). Protect

COMPARABLE PRODUCT

Ivomec® Pour-On, Coopers Paramax® Pour-On.

DOSING & ADMINISTRATION

application to cattle; do not use in other species. The formulation should be applied along the topline in a narrow strip extending from the

withers to the tailhead. The dose is 1 mL for each 10 kg of liveweight.

Dose the mob according to the heaviest animal by liveweight in each group (cows, bulls, calves, heifers etc.). When there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid overdosing. Do not underdose. Cattle should be weighed with either scales or with a weighband.

Liveweight (kg)	Dose (mL)
< 100	1 mL/ 10 kg
101 – 150	15
151 – 200	20
201 – 250	25
251 – 300	30
301 – 350	35
351 – 400	40
401 – 450	45
451 – 500	50
501 – 550	55
551 – 600	60
601 – 650	65

AVAILABLE SIZES 5 L, 20 L

For more information, visit pastoralag.com.au For product enquiries, email us at enquiries@pastoralag.com.au or call us on (03) 5820 8444

Coopers & Paramax are registered trade marks of Intervet Inc. a Delaware corporation. Ivomec is a registered trade mark of Boehringer Ingelheim Vetmedica GmbH.

*

INVALUABLE **WORM** CONTROL

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING FOR ANIMAL TREATMENT ONLY

PASTORAL / AG

FOR CATTLE

ACTIVE CONSTITUENT: 5 g/L IVERMECTIN







