## PASTORAL / AG

# TICKGARD POUR-ON

**ACTIVE CONSTITUENT: 25 g/L FLUAZURON** 

**FOR CATTLE** 

ALSO CONTAINS: 415 g/L 1-METHYL-2-PYRROLIDONE (solvent), 100 g/L 1-DODECYL-2-PYRROLIDONE (solvent), 100 g/L 1-OCTYL-2-PYRROLIDONE (solvent)

Pastoral Ag Tickgard is a ready to use pour-on formulation that contains the powerful insect growth regulator (IGR), fluazuron 25g/L, which interferes with chitin synthesis of the larval stages of the cattle tick. Controls susceptible strains of cattle tick (Boophilus microplus) & Inhibits tick development within three days. When treated with Pastoral Ag Tickgard, immature ticks are unable to molt to the next stage of their life cycle, and no larvae will hatch from eggs laid by treated females, breaking the tick lifecycle.

#### **APPLICATOR**

#### WITHHOLDING PERIODS

MEAT: DO NOT USE less than 42 days before MILK: DO NOT USE on cows which are producing or

ESI: DO NOT USE less than 42 days before slaughter for export.

#### **STORAGE**

### **COMPARABLE PRODUCTS**

Acatak™ Pour-On, OzTik Pour-On.

#### **DOSING & ADMINISTRATION**

but not before at least 6 weeks after the previous

ANIMALS & PESTS	STATES	BODYWEIGHT (kg)	DOSE (mL)
Beef cattle after weaning [Rhipicephalus (Boophilus) microplus]	QLD, NT, WA & NSW only	101 – 150	9
		151 – 200	12
		201 – 250	15
		251 – 300	
		301 – 350	21
		351 – 400	24
		401 – 450	27
		451 – 500	30
		501 – 550	33
		551 – 600	36
		601 – 650	39
Apply 3 mL for each 50 kg above 650 kg.			

**AVAILABLE SIZES** 5L | 20L

For more information, visit pastoralag.com.au For product enquiries, email us at enquiries@pastoralag.com.au or call us on (03) 5820 8444









CATTLE TICK **PROTECTION** 

REDUCE TICK PASTURE CONTAMINATION

PASTORAL AG POUR-ON



