

PASTORAL / AG™

DORALAB™ INJECTION FOR CATTLE AND PIGS

ACTIVE CONSTITUENT: 10 mg/mL DORAMECTIN

Doramectin is one of the most potent mectins and is trusted as an injectable for cattle due to its high efficacy combined with safety profile. PASTORAL AG DORALAB's HDPE plastic packaging prevents breakages in the yards.

PASTORAL AG DORALAB INJECTION has a broad spectrum of activity. It was effective in the treatment and control of the following species on cattle in trials conducted in Australia: Adult and immature Gastrointestinal Roundworms, Adult and immature Lungworm, Sucking Lice, Biting Lice, Cattle Tick, and Mange Mites.

Persistent Protection (Cattle):

Species	Period of protection following treatment
<i>Ostertagia ostertagi</i>	Up to 21 days
<i>Cooperia oncophora</i>	Up to 14 days
<i>Cooperia spp.</i>	Up to 21 days
<i>Haemonchus placei</i>	Up to 21 days
<i>Dictylocaulus viviparus</i>	Up to 28 days
<i>Oesophagostomum radiatum</i>	Up to 21 days
<i>Trichostrongylus axei</i>	Up to 21 days
<i>Bunostomum phlebotomum</i>	Up to 21 days

APPLICATOR

Cattle: PASTORAL AG 15ml or 20ml Variable Vaccinator.
Pigs: PASTORAL AG 2 or 5mL Variable Vaccinator.

WITHHOLDING PERIODS

MEAT:
CATTLE: 42 days.
PIGS: 35 days.
MILK: DO NOT USE in lactating cows or within 60 days of calving.
ESI: 42 days.

STORAGE

Store below 30°C (room temperature).

COMPARABLE PRODUCT

Zoetis Dectomax® Injectable Endectocide

For more information, visit pastoralag.com.au

For product enquiries, email us at enquiries@pastoralag.com.au or call us on (03) 5820 8444

Zoetis & Dectomx are registered trade marks of Zoetis Services LLC.

DOSING & ADMINISTRATION

CATTLE:

Administer 1 mL per 50 kg bodyweight subcutaneously (SC) (under the skin) into the neck region and preferably high up behind the ear. Use the dose rate, according to the weight of the animal, given in the table below.

Bodyweight (kg)	Dose (mL)
40 – 50	1
51 – 75	1.5
76 – 100	2
101 – 150	3
151 – 200	4
201 – 250	5
251 – 300	6
301 – 350	7
351 – 400	8
401 – 450	9
451 – 500	10
501 – 550	11
551 – 600	12
601 – 650	13

PIGS:

Administer 1 mL per 33 kg (0.3 mL per 10 kg) of bodyweight intramuscularly (IM) into the neck region, preferably high up behind the ear. A single treatment of 0.3 mL per 10 kg bodyweight (1 mL per 33 kg) should be given. Pigs should be weighed before treatment either with scales or a weighband. Piglets weighing 16 kg or less should be dosed in accordance with the following table.


Bodyweight (kg)	Dose (mL)
5 – 7	0.2
8 – 10	0.3
11 – 13	0.4
14 – 16	0.5

AVAILABLE SIZES

500ml

DORAMECTIN
POWER,

PLASTIC BOTTLE
CONVENIENCE



EXCLUSIVELY AVAILABLE FROM



MADE IN AUSTRALIA