

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

STARMEC®

IVERMECTIN INJECTION FOR CATTLE



ACTIVE CONSTITUENT: 10 mg/mL IVERMECTIN

For the treatment and control of ivermectin-sensitive internal and external parasites of cattle.

500 mL

Read the carton for full instructions.

DIRECTIONS FOR USE

Restraints:

DO NOT USE in lactating cows or within 28 days of calving where milk may be used or processed for human consumption.

Retreatment Interval: DO NOT re-treat less than 42 days after the last treatment.

Contraindications:

This product is contraindicated for use in calves at or less than 4 months of age.

Dosage and Administration:

Use the contents within 24 hours of first broaching of the vial.

Discard the unused portion.

Dose rate: 1 mL/50 kg liveweight SC. Refer to the dosage table on the carton as a guide.

WITHHOLDING PERIODS

MEAT: 28 days.

MILK: DO NOT USE in lactating cows or within 28 days of calving.

TRADE ADVICE ESI: 42 days.

SAFETY DIRECTIONS Poisonous if swallowed. Avoid contact with eyes, skin and clothing.

FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ENVIRONMENTAL PROTECTION Ivermectin is extremely toxic to aquatic species.

DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL Dispose of empty container by wrapping with paper and putting in garbage.

STORAGE Store below 25°C (air conditioning). Store bottle in carton to protect from light.

May damage fertility or the unborn child.

May cause harm to breast-fed children.

Do not breathe fumes, mists, vapours or spray.

Avoid contact during pregnancy or while nursing.

Wash contacted areas thoroughly after handling.

Do not eat, drink or smoke when using this product.

Use personal protective equipment as required.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

BATCH:

EXP: