

148mm

430mm

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

EXITRAZ EC

CATTLE AND PIG SPRAY

Active Constituent: 125 g/L Amitraz
Also contains: 600 g/L Hydrocarbon liquid (solvent)



For the control of: Organophosphate and Synthetic Pyrethroid resistant
Cattle Tick, Paralysis Tick and Bush Tick on Beef and Dairy Cattle, and
Cattle Tick on Sheep, Goats, Deer and Certain Circus Animals in NSW
only, and Mange on Pigs.



2 L



ZERO (0) DAYS
WITHHOLDING PERIODS
FOR MEAT AND MILK



DIRECTIONS FOR USE

Contraindications
This product should not be used on horses, other equines, felines or Chihuahua dogs.

Precautions
Treat animals in accordance with good agricultural practice. Animals should not be treated in extremely hot weather (>40°C) or if under heat stress. Ensure adequate shade is available post-treatment.

Dosage and Administration

LIVESTOCK	PEST	STATE	RATE	CRITICAL COMMENTS	
Cattle	Cattle Tick [<i>Rhipicephalus</i> (<i>Boophilus</i>) <i>microplus</i>]	NSW, QLD, WA and NT only	400 mL per 200 L water	Treat at intervals of 19-21 days	Controls organophosphate and synthetic pyrethroid resistant strains of the Cattle Tick. Refer also to General Directions
	Paralysis Tick (<i>Ixodes holocyclus</i>)				
	Bush Tick (NZ Cattle Tick) (<i>Haemaphysalis longicornis</i>)	NSW, QLD, WA, NT and VIC		Treat at intervals of 7-21 days	
Sheep, goats, deer and certain circus animals	Cattle Tick [<i>Rhipicephalus</i> (<i>Boophilus</i>) <i>microplus</i>]	NSW only	400 mL per 200 L water	Treat animals on quarantined holdings at 14 to 21 day intervals. Treat animals being moved at 3 to 7 day intervals.	
Pigs	Mange (<i>Sarcoptes scabiei</i> var. <i>suis</i>)	All States	40 mL per 10 L water	Remove feed and bedding from pens, cover drinking bowls and clean out pen. Spray pigs with minimum 2 L spray wash, especially inside ears and legs, under jowls and areas covered by scabs. Replace discarded bedding with clean material. Repeat treatment after 7-10 days. Refer also to General Directions.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

General Directions
Hand Spraying: Ensure all parts of the animal are thoroughly wet, especially belly, inside legs and ears (most important for pigs). Animals should be restrained in a crush for spraying. A pump capable of delivering a coarse spray should be used. Knapsack sprays are NOT recommended.

A fresh spray wash should be made up for each day's treatment. Clean out the spray reservoir at the end of each day and discard any remaining wash. Two treatments at 7-10 day intervals are recommended for sows and gilts before entering farrowing pens, for piglets at weaning and for pigs newly brought inside the piggery. Boars should be treated every 3 months.

Spray Races:

ACTION	UNSTABILISED	STABILISED
	Stabilisation is not required when all spray wash is used within 24 hours.	Stabilisation is required when unused wash is retained for later use.
Initial Charging	400 mL per 200 L water	
Mixing	Pour required amount of EXITRAZ EC into a bucket with approximately 10 L of water and stir thoroughly. Pour contents of bucket into reservoir or sump. Rinse bucket and add rinsings. Fill reservoir with total volume of water required.	
Stirring	Spraywash in sump must be stirred thoroughly at commencement of each spraying session and again if there is a break for more than half an hour during spraying.	
Reinforcing	Each time the level of spraywash in the sump drops by 200 L add 250 mL EXITRAZ EC, without adding water, and continue spraying. If level of wash is allowed to fall excessively without reinforcement, concentration of EXITRAZ EC will fall and efficiency will be reduced. Repeat reinforcement until no more wash can be pumped out. If continuing spraying, make up more wash at the initial charge rate and complete spraying. When wash is polluted clean out sump before recharging.	
Stabilising	Important: Because the wash is not stabilised, any remaining wash must be discarded and a new batch made up for the next spraying session. Alternatively, wash may be stabilised as opposite.	If, after spraying, unused wash is retained in the sump, add hydrated lime (EXITRAZ Stabiliser). Mix lime to a smooth paste with water and strain into wash at the rate of 2 kg per 200 L of spraywash and stir vigorously. Further additions of hydrated lime at the rate of 1 kg per 200 L wash must be made at intervals of 2 months when regular spraying is not in progress. Stir or agitate at regular intervals.
Sampling	Sampling NOT required.	Stir the sump vigorously before taking a sample from a nozzle on the race. Analysis of dip samples is available through the Biosecurity Queensland, Chemical Residue Laboratory, Health and Food Sciences Precinct, 39 Kessels Road, Coopers Plains, QLD 4108. Suitable sample bottles are obtainable from your local agent.

WITHHOLDING PERIODS

MEAT: Zero (0) days
MILK: Zero (0) days

TRADE ADVICE – EXPORT SLAUGHTER INTERVAL (ESI)
Cattle, Sheep: Zero (0) days. Before using this product, confirm the current ESI from The Hunter River Company on (03) 5821 0922 or the APVMA website (www.apvma.gov.au/residues).
Pigs, Goats and Deer: This product does not have an ESI established. For advice on the ESI, contact the The Hunter River Company on (03) 5821 0922 before using this product.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Do NOT inhale spray mist. When preparing spray or using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash contaminated clothing, gloves and face shield.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ADDITIONAL USER SAFETY INFORMATION

Additional information is listed in the safety data sheet.

ENVIRONMENTAL PROTECTION

EXITRAZ EC is toxic to fish. Do NOT contaminate dams, rivers or streams with EXITRAZ EC, used containers or dip or spray wash containing EXITRAZ EC. DO NOT use empty containers for any other purpose. If used in accordance with the directions and recommendations, EXITRAZ EC will not give rise to hazards to livestock or wildlife.

DISPOSAL

Triple-rinse container into the medicated spray tank. Do not dispose of undiluted chemicals on-site. If the container has the drum MUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (room temperature) in the closed original container in a dry, well-ventilated area, out of direct sunlight.

Harmful if swallowed.
May be fatal if swallowed and enters airways.
May cause an allergic skin reaction.
Use only outdoors or in a well ventilated area.
Wear protective gloves, protective clothing and eye or face protection.
Take off contaminated clothing and wash before reuse.
IF SWALLOWED: Immediately call a POISON CENTRE or doctor. Rinse mouth. Do NOT induce vomiting.
IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.
Collect spillage.

B:

EXP:

The Hunter River Company Pty Limited
ACN 133 798 615 | 74-76 Drummond Rd,
Shepparton VIC 3630 | Ph: (03) 5821 0922

APVMA Approval No.: 85136/129770

240713L-3