

SURESHOT™

DUO COMBINATION

POUR-ON FOR CATTLE



ACTIVE CONSTITUENTS:

200 mg/mL LEVAMISOLE, 10 mg/mL ABAMECTIN

Also contains: 540 mg/mL N-METHYL-2-PYRROLIDONE,
250 mg/mL DIETHYLENE GLYCOL MONOBUTYL ETHER (solvent)



1L 2.5L 5L

For the treatment and control of roundworms, including macrocyclic lactone or levamisole resistant strains, and external parasites in cattle.

The combined power of abamectin and levamisole in a pour-on formulation provide control over parasites resistant to mectins or BZs, as well as slowing the onset of parasite resistance on the property. Sureshot Duo Combination is a convenient low dose 1ml/20kg pour-on formulation that not only controls internal parasites, will also control ecto parasites such as lice, buffalo fly and cattle tick.

APPLICATOR

30ml Variable Applicator – Pour-On Nozzle.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 56 days before slaughter for human consumption. Calves MUST NOT be slaughtered for human consumption within 60 days of treatment of dams.
MILK: DO NOT USE in lactating cows or within 60 days of calving where milk may be used or processed for human consumption. If cows calve earlier than 60 days after treatment, milk may contain residues. This milk MUST NOT be used for human consumption, supplied for processing or fed to bobby calves for at least 60 days following treatment.
ESI: DO NOT USE less than 70 days before slaughter for export.

STORAGE

Store below 25°C (air conditioning). Protect from light.

DOSING & ADMINISTRATION

SURESHOT DUO POUR-ON dose rate is 1 mL/20 kg body weight (0.5 mg/kg abamectin and 10 mg/kg levamisole). Apply along the backline of the animal, in a strip starting between the shoulder blades.

LIVEWEIGHT (KG)	DOSE VOLUME (ML)
51-100	5
101-150	7.5
151-200	10
201-250	12.5
251-300	15
301-350	17.5
351-400	20
401-450	22.5
451-500	25
501-550	27.5
551-600	30
601-650	32.5

AVAILABLE SIZES

1L, 2.5L, 5L

For more information, visit hunterriverco.com.au

