

FRONT

145mm



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

SURESHOT™
DUO COMBINATION
POUR-ON FOR CATTLE

ACTIVE CONSTITUENTS: 200 mg/mL LEVAMISOLE, 10 mg/mL ABAMECTIN
Also contains: 540 mg/mL N-METHYL-2-PYRROLIDONE,
250 mg/mL DIETHYLENE GLYCOL MONOBUTYL ETHER (solvent)

For the treatment and control of roundworms, including macrocyclic lactone or levamisole resistant strains, and external parasites in cattle.

1L



31423F-3

HUNTER RIVER CO.

109mm

BACK

SURESHOT DUO POUR-ON contains abamectin, a member of the macrocyclic lactone family of chemicals. It is effective against sensitive strains of internal and external parasites. It also contains levamisole, a member of the imidazothiazole family of chemicals, which is effective against roundworms, including lungworm. Resistance may develop to any chemical.

Read the carton for full instructions.

DIRECTIONS FOR USE

Restrictions:

DO NOT USE in lactating cows or within 60 days of calving where milk may be used or processed for human consumption. **DO NOT USE** in calves under 50 kg bodyweight. **Re-treatment interval: DO NOT re-treat animals for 35 days after last treatment.**

Dosage and Administration:

SURESHOT DUO POUR-ON dose rate is 1 mL/20 kg body weight (0.5 mg/kg abamectin and 10 mg/kg levamisole). Apply along the backline of the animal, using the SURESHOT DUO POUR-ON applicator, in a strip starting between the shoulder blades. This product is not to be given

orally or injected. Only use the pour-on applicator supplied for use with SURESHOT DUO POUR-ON. SURESHOT DUO POUR-ON is not compatible with other guns. After use, rinse gun with warm soapy water.

Refer to the carton for dosage table.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 56 days before slaughter. Calves MUST NOT be slaughtered for human consumption within 60 days of treatment of dams. MILK: DO NOT USE in lactating cows or within 60 days of calving. If cows calve earlier than 60 days after treatment, milk may contain residues. This milk MUST NOT be used for human consumption, supplied for processing or fed to bobby calves for at least 60 days following treatment.

TRADE ADVICE
EXPORT SLAUGHTER INTERVAL (ESI): 70 days.

SAFETY DIRECTIONS

Harmful if swallowed or absorbed by skin contact. Repeated exposure may cause allergic disorders. Will damage the eyes and skin. Avoid contact with eyes and skin. If clothing becomes contaminated with product, remove clothing immediately. If product is in eyes, wash it out immediately with water. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat, elbow-length chemical resistant gloves and goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

ENVIRONMENTAL PROTECTION

Abamectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Dispose of container by wrapping with paper and putting in garbage.

STORAGE

Store below 25°C (air conditioning). Protect from light.

Fatal if swallowed.
Toxic if inhaled.
May cause respiratory irritation.
May damage the unborn child.
May cause damage to organs through prolonged or repeated exposure.

Wash all exposed external body areas thoroughly after handling.
Do not eat, drink or smoke when using this product.
Use only outdoors or in a well-ventilated area.
IF INHALED: Remove person to fresh air and keep comfortable for breathing.

BATCH:

EXP:

31423B-4

The Hunter River Company Pty Limited ACN 133 798 615 | 74-76 Drummond Rd, Shepparton VIC 3630 | Ph: (03) 5820 8400
APVMA Approval No.: 87741/129840

141mm

104mm