


FRONT



LEFT SIDE

ABASTAR PLUS Selenium LV contains abamectin, a member of the avermectin family of drenches. It is effective against abamectin sensitive strains of the following internal parasites:

Adult and immature gastrointestinal roundworms:
Haemonchus contortus – Barber's pole worm
Ostertagia spp. – Small brown stomach worms
Trichostrongylus spp. – Black scour worms
Cooperia spp. – Small intestinal worms
Nematodirus spp. – Thin-necked intestinal worms
Oesophagostomum venulosum – Large bowel worm

Lungworm:
Dictyocaulus filaria

For the control of **Sheep Nasal bot** – *Oestrus ovis*.

For the treatment and prevention of selenium deficiencies in sheep.

Resistance may develop to any chemical.

DIRECTIONS FOR USE:

Restrictions:
DO NOT USE in lactating ewes where milk may be used or processed for human consumption.

Contraindications:
 This product is contraindicated for use in lambs under 6 weeks of age or less than 10 kg bodyweight.
 This product is contraindicated for use when selenium intake from pasture is high or selenium is provided by other means (such as top dressing, vaccine, pellets or selenium drenches) if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian.

Precautions:
 Use with caution during early pregnancy, accurate dosing is important and particular care should be taken to avoid overdosing. This product should only be used when drenching is required according to a worm control program. If selenium is required at other times it should be administered by way of other products.

Dosage and Administration:
 1 mL ABASTAR PLUS Selenium LV per 10 kg bodyweight. Drench sheep orally, using an appropriate drench gun.

Dose rate table:

BODYWEIGHT (KG)	DOSE (ML)	NO. OF SHEEP TREATED
16 – 20	2.0	5000
21 – 25	2.5	4000
26 – 30	3.0	3333
31 – 35	3.5	2857
36 – 40	4.0	2500
41 – 45	4.5	2222
46 – 50	5.0	2000
51 – 55	5.5	1818
56 – 60	6.0	1666
61 – 65	6.5	1538
66 – 70	7.0	1428
71 – 75	7.5	1333

Animals heavier than 75 kg to be dosed at 1 mL per 10 kg.

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

9 348896 000508

24703LEFT-2

100mm

150mm

RIGHT SIDE

General Directions:
 To control white muscle disease in lambs, treat pregnant ewes 2-6 weeks before lambing and at marking. On properties where selenium responsive illthrift has been diagnosed, treat lambs at weaning and again 6-10 weeks later. During drenching, regularly check drench gun to ensure it is delivering the correct dose. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your local area to reduce development of resistance. It is advisable that a drench resistance test be conducted before any drench is used.

WITHHOLDING PERIODS
MEAT: DO NOT USE less than 14 days before slaughter for human consumption.
MILK: DO NOT USE in lactating ewes where milk may be used or processed for human consumption.

TRADE ADVICE
EXPORT SLAUGHTER INTERVAL (ESI): An ESI has not been established for this product. Note—observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from The Hunter River Company on (03) 5820 8400 before using this product.

SAFETY DIRECTIONS
 Poisonous if absorbed by skin contact or swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and using the product, wear rubber gloves. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID INSTRUCTIONS
 If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ADDITIONAL USER SAFETY
 Additional information is listed in the safety data sheet.

ENVIRONMENTAL PROTECTION
 Abamectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL
 Triple-rinse container and dispose of rinse in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately at the drumMUSTER collection site. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE
 Store below 30°C (room temperature) in original container, tightly closed in a cool, dry place. Protect from light.

Harmful if inhaled.
 Avoid breathing fumes, mists, vapours or spray.
 Do not eat, drink or smoke when using this product.
 Use only outdoors or in a well ventilated area.
 Use personal protective equipment as required.
IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.
IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.
 Not combustible. Use extinguishing media suited to burning materials.
 Store in a well-ventilated place.

BATCH:
 EXP:

24703RIGHT-2

24703F-2